



PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	MIM	Books	
Search	PubMed	▼	for					Go	Clear
Limits    Preview/Index    History    Clipboard    Details									

Display	Abstract	▼	Show:	20	▼	Sort	▼	Send to	Text	▼
---------	----------	---	-------	----	---	------	---	---------	------	---

☐ 1: Br J Cancer 1992 Sep;66(3):474-8

[Related Articles, Links](#)

Entrez  
PubMed

## **A monoclonal antibody-beta-glucuronidase conjugate as activator of the prodrug epirubicin-glucuronide for specific treatment of cancer.**

**Haisma HJ, Boven E, van Muijen M, de Jong J, van der Vijgh WJ, Pinedo HM.**

PubMed  
Services

Department of Medical Oncology, Free University Hospital, Amsterdam, The Netherlands.

Related  
Resources

The anti-pan carcinoma monoclonal antibody (MAb) 323/A3, linked to E. coli-derived beta-glucuronidase (GUS) was used to study the tumour-site-selective activation of the prodrug Epirubicin-glucuronide (Epi-glu). Epi-glu was isolated from the urine of patients treated with Epirubicin (Epi) by reversed phase chromatography on a silica-C18 column. Epi-glu was stable in human blood and was not converted into Epi by A2780, MCF-7, or OVCAR-3 cancer cells, despite the presence of intracellular GUS. The stability of the prodrug was confirmed in BALB/c mice. MAb 323/A3 and GUS were linked through a stable thioether bond. The conjugate (1:1) was purified by ion exchange and gel filtration chromatography. Binding to target cells revealed an immunoreactivity of at least 60% and good retention of enzyme activity. A protein dye (sulforhodamine B) assay was used to analyse cytotoxicity. Epi (IC<sub>50</sub> of 0.003-0.2 microM) was 100-1,000 times more toxic than Epi-glu (IC<sub>50</sub> of greater than 20 microM), when cancer cells were exposed for 4 or 24 h to the drugs. The low cytotoxicity of Epi-glu was most likely due to the reduced cellular uptake rate of the prodrug (2.7 pmol 10<sup>-6</sup> cells min<sup>-1</sup>) as compared to that of the parent compound (25 pmol 10<sup>-6</sup> cells min<sup>-1</sup>). Pretreatment of antigen-positive cells with the 323/A3-GUS conjugate prior to prodrug exposure completely restored cytotoxicity as a result from hydrolysis of Epi-glu into Epi. Our results demonstrate that the 323/A3-GUS conjugate can specifically activate the stable non-toxic prodrug Epi-glu at the tumour cell level.

PMID: 1520585 [PubMed - indexed for MEDLINE]

Display	Abstract	▼	Show:	20	▼	Sort	▼	Send to	Text	▼
---------	----------	---	-------	----	---	------	---	---------	------	---

[Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)  
[Department of Health & Human Services](#)

Freedom of Information Act | Disclaimer

March 22, 2003 10:59 AM

■